

Cystic Fibrosis and bone health

This factsheet has been written to assist you and your medical advisors. It is not intended to replace any advice you may receive from your Specialist CF Centre or Clinic.

Introduction

As we get older our bones become thinner and weaker, and may become more susceptible to fracture. However in people with Cystic Fibrosis, thinning of the bones can occur at an earlier age. This factsheet describes how bone health can be affected in CF and the implications of this, how bone strength is measured, and options for prevention and treatment of thinning bones.

This factsheet was written by Dr Steve Conway, Clinical Director of CF Services, St James's University Hospital, Leeds and members of the Cystic Fibrosis Trust Bone Mineralisation Working Group.

Last reviewed **January 2011**

www.cftrust.org.uk

Patron: HRH Princess Alexandra, the Hon. Lady Ogilvy, KG, GCVO
Cystic Fibrosis Trust registered as a charity in England and Wales (1079049) and in Scotland (SC040196)
A company limited by guarantee registered in England and Wales number 3880213
Registered office: 11 London Road, Bromley, Kent BR1 1BY

Osteopenia and osteoporosis

Osteopenia and osteoporosis are terms that mean bones are “thinner” than they should be. A more scientific description is that the skeleton has a low bone mass. This means that the bones are more fragile and more likely to fracture.

What is Bone Mineral Density (BMD)?

Bone mineral density (BMD) is a measure of bone mineralisation – the level of minerals contained in bone. The measurements are an indicator of the bone mass and therefore how strong the bones are.

What are Bone Mineral Density “Z” and “T” scores?

BMD measurements are scored according to how much they fall below or above the population average value. BMD results can be reported as Z-scores or T-scores, but Z-scores are usually the most appropriate method in people with Cystic Fibrosis. The Z-score compares a measured BMD value to the average value obtained from a healthy population of the same sex and age as the patient.

Generally speaking, BMD is considered low in people with CF when the BMD Z-score in the spine or hip is lower than -2 . The T-score compares the measured BMD value to the average value obtained from a healthy young adult population. (In young adult life our BMD is at its highest value.)

What happens in normal bones?

- The skeleton provides a rigid framework, a site for attachment of muscles, protects internal organs, and is important in maintaining normal body calcium and phosphate levels.
- Large amounts of bone minerals are acquired in childhood and especially in adolescence during the pubertal growth spurt. About 90% of adult bone mineral is acquired by late adolescence with peak bone mass reached

between 20 to 30 years of age. We usually start to lose bone from about 40 years of age.

- Old bone is always being removed (bone resorption) and replaced by new bone (bone formation). About 10% of adult bone is replaced by this process each year.

Lifetime changes in bone mass



What about in CF?

People with CF are at risk of early thinning of the bones. BMD is usually normal in well-nourished children with CF who have good lung function. Even so many patients fail to gain bone normally, or experience premature bone loss, in adolescence. About one third of adults with CF have low bone mineral density which may predispose them to bone fractures.

How is BMD measured?

BMD is usually measured by dual energy X-ray absorptiometry (DXA).

Dual energy X-ray absorptiometry

Dual energy X-ray absorptiometry (DXA) is used to measure BMD in the lumbar spine (below the chest and above the pelvis), the top of the leg (hip), the wrist, and the whole body. The scan at each site takes approximately two to three minutes and involves a low radiation dose, less than the dose of naturally occurring background radiation and much less than that needed for a chest x-ray. The BMD is scored and Z and T values calculated. It is important to note that the BMD score may not accurately predict the fracture risk in people with Cystic Fibrosis.

A DXA scan should be performed from about ten years of age and repeated every one to three years, the frequency determined by clinical need. This should be organised by your CF Centre and the results interpreted in liaison with the DXA scanning department.

What causes low bone mineral density in CF?

CF-related low BMD probably has many causes:

- 1) Bone health is strongly linked to the severity of lung disease and body weight. Almost all severely ill patients have low BMD. Conversely patients with normal weight and height, and good lung function, have near normal bone density. Lung infection in CF is associated with more intense inflammation and raised levels of chemicals called cytokines in blood and sputum. These cytokines increase the activity of cells which break down bone.
- 2) Apart from the general malnutrition associated with more severe disease, many aspects of nutrition affect bone status, including vitamins (e.g., vitamins D and K), minerals (e.g., calcium), and protein intake.
- 3) Steroids cause rapid bone loss, especially during the first year of therapy. They reduce calcium absorption from the gut, increase calcium loss in the urine, decrease the number of bone forming cells and increase bone resorption. Most of the larger studies in individuals with CF have shown an association between oral steroid use and low BMD.
- 4) A number of studies in people who do not have CF have found an association between greater physical activity and higher BMD levels. However, it is not known if weight-bearing exercise in people with CF can increase peak bone mass, preserve BMD or increase BMD in those with low BMD.

- 5) Despite most children with CF achieving normal or near normal growth, puberty is often delayed. Bone mineral deficiencies resulting from late onset of puberty may not be corrected when puberty eventually starts. Low sex hormone levels, oestrogen and testosterone, are associated with low BMD in adults with Cystic Fibrosis.
- 6) Low bone mineral density has been associated with Cystic Fibrosis related diabetes.
- 7) There may be a direct link between CF related low BMD and the abnormal protein produced by the CF gene.

Why is it important to prevent fractures?

Fractures in the spine and ribs are painful and therefore may interfere with effective physiotherapy and airway clearance, resulting in a decline in lung function.

How can low BMD be prevented and treated?

- People with CF should be screened for low BMD by regular DXA scans and chest X-rays. The latter should be examined specifically for vertebral fractures. Risk factors that we can treat such as low vitamin D or calcium levels should be monitored and low levels corrected with extra dietary supplements.
- Lung infection damages bone health. Therefore treatments that prevent the progression of lung disease should be optimised.
- Optimising nutrition through dietary intervention, and specialist dietetic supervision of appropriate and proactive interventions with oral supplements and nasogastric or gastrostomy tube feeds is fundamental to care. Vitamin D and vitamin K status and calcium intake should be optimised through

frequent contact with a specialist CF dietitian. Vitamin D and calcium levels should be measured at least annually. There is not yet sufficient evidence to recommend universal vitamin K supplementation for bone health in CF, but supplementation should be considered in certain clinical situations e.g., for patients with a low BMD, or liver disease. Patients should have access to a specialist dietitian at each admission, outpatient review, and annual assessment.

- Weight bearing physical activity should be encouraged.
- A specialist CF physiotherapist should develop an exercise programme where appropriate for an individual patient's abilities and needs.
- Glucocorticoid treatment (prednisolone and inhaled steroids) should be kept to a minimum.
- Delayed puberty and low blood levels of testosterone and oestrogens should be looked for and treated.
- As long acting "depot" and possibly other progesterone-only contraception preparations may reduce BMD, particularly in adolescents, alternative contraceptive preparations may be suggested.
- Smoking and alcohol can have damaging effects on bone health and should be avoided.

Are there any specific drug treatments?

- Bisphosphonates (e.g., Aledronate, Residronate) are a family of drugs that reduce bone breakdown and may directly stimulate the cells that manufacture bone. Non-CF studies have shown benefit in the treatment of postmenopausal osteoporosis and there is increasing evidence for their efficacy in adults

with Cystic Fibrosis.

- Bisphosphonates should not be used during pregnancy and should therefore be used with caution in young women.
- Bisphosphonates are best absorbed in the absence of food and should be taken in the fasting state. This may be difficult for people with CFRD or for those who feed overnight. Calcium supplements impair the absorption of bisphosphonates and the two should not be taken together.
- Bisphosphonates can cause ulceration when passing from the mouth to the stomach so swallowing should be helped by patients staying upright for about 30 minutes after taking the drug.
- Bisphosphonates are not licensed for use in children and although they appear to be relatively safe even when used for long periods in other paediatric bone disease, experience is limited.
- We recommend that treatment of children with bisphosphonates is supervised by a major CF Centre or by a paediatric bone specialist.

Further reading

The Cystic Fibrosis Trust has published a consensus document entitled Bone Mineralisation in Cystic Fibrosis. This can be downloaded from our website www.cftrust.org.uk or hard copies obtained by contacting the Cystic Fibrosis Trust.

The following websites may also be useful:

National Osteoporosis Society www.nos.org.uk

National Osteoporosis Foundation www.nof.org

National Institute of Health www.osteop.org

NHS Direct Online www.nhsdirect.nhs.uk

International Osteoporosis Foundation www.osteofound.org

Cystic Fibrosis Medicine www.cysticfibrosismedicine.com

Further information

For further information about any aspect of Cystic Fibrosis, please contact:

Cystic Fibrosis Trust
11 London Road
Bromley
Kent BR1 1BY
T 020 8464 7211
www.cftrust.org.uk
enquiries@cftrust.org.uk

Helplines

General helpline 0300 373 1000

Welfare Grants helpline 0300 373 1020

Benefits helpline 0300 373 1050

Other publications

The Cystic Fibrosis Trust publishes a range of publications on a wide variety of topics relating to Cystic Fibrosis. You can view our complete publications list, order publications and download publications free of charge from the CF Trust website www.cftrust.org.uk/publications. Alternatively, you can telephone the CF Trust switchboard on 020 8464 7211.

We would welcome your feedback on this or any other Cystic Fibrosis Trust publications. Please email publications@cftrust.org.uk.

© Cystic Fibrosis Trust 2011. This factsheet may be copied in whole or in part, without prior permission being sought from the copyright holder, provided the purpose of copying is not for commercial gain and due acknowledgement is given.